



# Fixed point results on metric and partial metric spaces via simulation functions

Antonella Nastasi, Pasquale Vetro\*

*Dipartimento di Matematica e Informatica, Università degli Studi di Palermo, via Archirafi 34, 90123 Palermo, Italy.*

---

## Abstract

We prove existence and uniqueness of fixed point, by using a simulation function and a lower semi-continuous function in the setting of metric space. As consequences of this study, we deduce several related fixed point results, in metric and partial metric spaces. An example is given to support the new theory. ©2015 All rights reserved.

*Keywords:* Fixed point, metric space, partial metric space, nonlinear contraction, simulation function.  
*2010 MSC:* 47H10, 54H25.

---

## 1. Introduction

Metric fixed point theory is a vivid topic, which furnishes useful methods and notions for dealing with various problems. In particular, we refer to the existence of solutions of mathematical problems reducible to equivalent fixed point problems. Thus we recall that Banach contraction principle [2] is at the foundation of this theory. However, the potentiality of fixed point approaches attracted many scientists and hence there is a wide literature available for interested readers, see for instance [3, 7, 8, 9, 11, 18, 20, 21, 24, 25, 27]. We give some details on the notions and ideas used in this study.

First, the notion of partial metric space was introduced in 1994 by Matthews [12] as a part of the study of denotational semantics of data for networks. Clearly, this setting is a generalization of the classical concept of metric space. Also, some authors discussed the existence of several connections between partial metrics and topological aspects of domain theory, see for instance [5, 6, 13, 17].

Second, the notion of  $\mathcal{Z}$ -contraction was introduced in 2014 by Khojasteh *et al.* [10]. This concept is a new type of nonlinear contraction defined by using a specific simulation function. Of course, they proved existence and uniqueness of fixed points for  $\mathcal{Z}$ -contraction mappings. We point out that the advantage of

---

\*Corresponding author

*Email addresses:* [ella.nastasi.93@gmail.com](mailto:ella.nastasi.93@gmail.com) (Antonella Nastasi), [pasquale.vetro@unipa.it](mailto:pasquale.vetro@unipa.it) (Pasquale Vetro)

this method is in providing a unique point of view for several fixed point problems; see recent results in [1, 23].

Finally, Samet *et al.* [26], and Vetro and Vetro [28] discussed fixed point results by using semi-continuous functions in metric spaces that generalize and improve many existing fixed point theorems in the literature. As an application of presented results, the authors gave some theorems in the setting of partial metric spaces. In this paper, we use the ideas in [26, 28] and the notion of simulation function to establish existence and uniqueness of fixed points. As consequences of this study, we deduce several related fixed point results, in metric and partial metric spaces. Also, an example is given to support the new theory.

## 2. Preliminaries

In this section we recall some definitions and some properties of partial metric spaces that can be found in [12, 13, 14, 15, 16]. We start with a definition. Precisely, a partial metric on a non-empty set  $X$  is a function  $p : X \times X \rightarrow [0, +\infty[$  such that, for all  $x, y, z \in X$ , we have

$$(p_1) \quad x = y \Leftrightarrow p(x, x) = p(x, y) = p(y, y),$$

$$(p_2) \quad p(x, x) \leq p(x, y),$$

$$(p_3) \quad p(x, y) = p(y, x),$$

$$(p_4) \quad p(x, y) \leq p(x, z) + p(z, y) - p(z, z).$$

Then, a partial metric space is a pair  $(X, p)$ , where  $X$  is a non-empty set and  $p$  is a partial metric on  $X$ . Notice that, if  $p(x, y) = 0$ , then from  $(p_1)$  and  $(p_2)$  it follows that  $x = y$ . However, if  $x = y$ , then  $p(x, y)$  may not be 0. A classic example of partial metric space is the pair  $([0, +\infty[, p)$ , where  $p(x, y) = \max\{x, y\}$  for all  $x, y \in [0, +\infty[$ , see also [12].

Every partial metric  $p : X \times X \rightarrow [0, +\infty[$  generates a  $T_0$  topology  $\tau_p$  on  $X$  which has as a base the family of open  $p$ -balls  $\{B_p(x, \varepsilon) : x \in X, \varepsilon > 0\}$ , where

$$B_p(x, \varepsilon) = \{y \in X : p(x, y) < p(x, x) + \varepsilon\}$$

for all  $x \in X$  and  $\varepsilon > 0$ .

Let  $(X, p)$  be a partial metric space. A sequence  $\{x_n\}$  in  $(X, p)$  converges to a point  $x \in X$  if and only if  $p(x, x) = \lim_{n \rightarrow +\infty} p(x, x_n)$ . A sequence  $\{x_n\}$  in  $(X, p)$  is called a Cauchy sequence if there exists (and is finite)  $\lim_{n, m \rightarrow +\infty} p(x_n, x_m)$ .

A partial metric space  $(X, p)$  is said to be complete if every Cauchy sequence  $\{x_n\}$  in  $X$  converges, with respect to  $\tau_p$ , to a point  $x \in X$  such that  $p(x, x) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m)$ .

It is easy to see that, every closed subset of a complete partial metric space is complete.

Notice that if  $p$  is a partial metric on  $X$ , then the function  $p^s : X \times X \rightarrow [0, +\infty[$  given by

$$p^s(x, y) = 2p(x, y) - p(x, x) - p(y, y) \tag{2.1}$$

is a metric on  $X$ . Furthermore,  $\lim_{n \rightarrow +\infty} p^s(x_n, x) = 0$  if and only if

$$p(x, x) = \lim_{n \rightarrow +\infty} p(x_n, x) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m).$$

**Lemma 2.1.** *Let  $(X, p)$  be a partial metric space and let  $\varphi : X \rightarrow [0, +\infty[$  be defined by  $\varphi(x) = p(x, x)$  for all  $x \in X$ . Then the function  $\varphi$  is lower semi-continuous in the metric space  $(X, p^s)$ .*

*Proof.* Let  $\{x_n\} \subset X$  be a sequence which converges to  $x \in X$  in the metric space  $(X, p^s)$ , then

$$p(x, x) = \lim_{n \rightarrow +\infty} p(x_n, x_n) = \liminf_{n \rightarrow +\infty} p(x_n, x_n).$$

It follows that  $\varphi$  is lower semi-continuous in  $x$  and hence in  $X$ . □

**Lemma 2.2** ([12, 14]). *Let  $(X, p)$  be a partial metric space. Then*

- (a)  $\{x_n\}$  is a Cauchy sequence in  $(X, p)$  if and only if it is a Cauchy sequence in the metric space  $(X, p^s)$ .
- (b) A partial metric space  $(X, p)$  is complete if and only if the metric space  $(X, p^s)$  is complete.

Khojasteh *et al.* gave the following definition of simulation function, see [10].

**Definition 2.3.** A simulation function is a mapping  $\zeta : [0, +\infty[ \times [0, +\infty[ \rightarrow \mathbb{R}$  satisfying the following conditions:

- ( $\zeta_1$ )  $\zeta(0, 0) = 0$ ;
- ( $\zeta_2$ )  $\zeta(t, s) < s - t$ , for all  $t, s > 0$ ;
- ( $\zeta_3$ ) if  $\{t_n\}, \{s_n\}$  are sequences in  $]0, +\infty[$  such that  $\lim_{n \rightarrow +\infty} t_n = \lim_{n \rightarrow +\infty} s_n = \ell \in ]0, +\infty[$ , then

$$\limsup_{n \rightarrow +\infty} \zeta(t_n, s_n) < 0.$$

Consequently, they proved the following theorem.

**Theorem 2.4** ([10]). *Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow X$  be a  $\mathcal{Z}$ -contraction with respect to a certain simulation function  $\zeta$ , that is,*

$$\zeta(d(Tx, Ty), d(x, y)) \geq 0, \quad \text{for all } x, y \in X. \tag{2.2}$$

*Then  $T$  has a unique fixed point. Moreover, for every  $x_0 \in X$ , the Picard sequence  $\{T^n x_0\}$  converges to this fixed point.*

In [1], Argoubi *et al.* note that the condition ( $\zeta_1$ ) was not used for the proof of Theorem 2.4. Also they observed that taking  $x = y$  in (2.2), one has  $\zeta(0, 0) \geq 0$  and hence, if  $\zeta(0, 0) < 0$ , the set of operators  $T : X \rightarrow X$  satisfying (2.2) is empty.

Therefore, Argoubi *et al.* slightly revised the previous definition, by withdrawing the condition ( $\zeta_1$ ). For the sake of clarity, we consider the following definition.

**Definition 2.5.** A simulation function is a mapping  $\zeta : [0, +\infty[ \times [0, +\infty[ \rightarrow \mathbb{R}$  satisfying the conditions ( $\zeta_2$ ) and ( $\zeta_3$ ).

Note that every simulation function in (original) Khojasteh *et al.* sense (Definition 2.3) is a simulation function in Argoubi *et al.* sense (Definition 2.5), but the converse is not true; see example below.

**Example 2.6** ([1], Example 2.4). Let  $\zeta_\lambda : [0, +\infty[ \times [0, +\infty[ \rightarrow \mathbb{R}$  be the function defined by

$$\zeta_\lambda(t, s) = \begin{cases} 1 & \text{if } (s, t) = (0, 0), \\ \lambda s - t & \text{otherwise,} \end{cases}$$

where  $\lambda \in ]0, 1[$ . Then  $\zeta_\lambda$  satisfies ( $\zeta_2$ ) and ( $\zeta_3$ ) with  $\zeta_\lambda(0, 0) > 0$ .

For completeness we give some of the most interesting examples of simulation functions.

**Example 2.7.** Let  $\zeta : [0, +\infty[ \times [0, +\infty[ \rightarrow \mathbb{R}$ , be defined by

- (i)  $\zeta(t, s) = \lambda s - t$  for all  $t, s \in [0, +\infty[$ , where  $\lambda \in [0, 1[$ .

- (ii)  $\zeta(t, s) = \psi(s) - \varphi(t)$  for all  $t, s \in [0, +\infty[$ , where  $\varphi, \psi : [0, +\infty[ \rightarrow [0, +\infty[$  are two continuous functions such that  $\psi(t) = \varphi(t) = 0$  if and only if  $t = 0$  and  $\psi(t) < t \leq \varphi(t)$  for all  $t > 0$ .
- (iii)  $\zeta(t, s) = s - \frac{f(t, s)}{g(t, s)}t$  for all  $t, s \in [0, +\infty[$ , where  $f, g : [0, +\infty[ \times [0, +\infty[ \rightarrow [0, +\infty[$  are two continuous functions with respect to each variable such that  $f(t, s) > g(t, s)$  for all  $t, s > 0$ .
- (iv)  $\zeta(t, s) = s - \psi(s) - t$  for all  $t, s \in [0, +\infty[$ , where  $\psi : [0, +\infty[ \rightarrow [0, +\infty[$  is a lower semi-continuous function such that  $\psi(t) = 0$  if and only if  $t = 0$ .
- (v)  $\zeta(t, s) = s\psi(s) - t$  for all  $t, s \in [0, +\infty[$ , where  $\psi : [0, +\infty[ \rightarrow [0, 1[$  is such that  $\lim_{t \rightarrow r^+} \psi(t) < 1$  for all  $r > 0$ .

### 3. Fixed points via simulation functions

We start this section with the following auxiliary lemma. Also, we use simulation function in the sense of Definition 2.5. Let  $X \neq \emptyset$ ,  $T : X \rightarrow X$ ,  $x_0 \in X$  and  $x_n = Tx_{n-1}$  for all  $n \in \mathbb{N}$ . Then  $\{x_n\}$  is called sequence of Picard of initial point at  $x_0$ .

**Lemma 3.1.** *Let  $(X, d)$  be a metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist a simulation function  $\zeta$  and a function  $\varphi : X \rightarrow [0, +\infty[$  such that*

$$\zeta(d(Tx, Ty) + \varphi(Tx) + \varphi(Ty), d(x, y) + \varphi(x) + \varphi(y)) \geq 0 \quad \text{for all } x, y \in X. \tag{3.1}$$

Let  $\{x_n\}$  be a sequence of Picard of initial point at  $x_0 \in X$ . Suppose that  $x_{n-1} \neq x_n$  for all  $n \in \mathbb{N}$ . Then  $\{x_n\}$  is a Cauchy sequence.

*Proof.* Let  $x_0$  be an arbitrary point in  $X$  and let  $\{x_n\}$  be a sequence of Picard of initial point at  $x_0 \in X$ . We shall prove the lemma in three steps. The first step is to prove that

$$\lim_{n \rightarrow +\infty} d(x_{n-1}, x_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} \varphi(x_n) = 0. \tag{3.2}$$

From,  $x_{n-1} \neq x_n$  for all  $n \in \mathbb{N}$ , we deduce that

$$d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n) > 0 \quad \text{for all } n \in \mathbb{N}.$$

Using (3.1) and  $(\zeta_2)$ , with  $x = x_{n-1}$  and  $y = x_n$ , we have

$$\begin{aligned} 0 &\leq \zeta(d(x_n, x_{n+1}) + \varphi(x_n) + \varphi(x_{n+1}), d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n)) \\ &< d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n) - [d(x_n, x_{n+1}) + \varphi(x_n) + \varphi(x_{n+1})] \end{aligned}$$

for all  $n \in \mathbb{N}$ . The above inequality shows that

$$d(x_n, x_{n+1}) + \varphi(x_n) + \varphi(x_{n+1}) < d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n), \quad \text{for all } n \in \mathbb{N},$$

which implies that  $\{d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n)\}$  is a decreasing sequence of positive real numbers. Thus, there is some  $r \geq 0$  such that

$$\lim_{n \rightarrow +\infty} d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n) = r. \tag{3.3}$$

Suppose that  $r > 0$ . It follows from the condition  $(\zeta_3)$ , with  $t_n = d(x_n, x_{n+1}) + \varphi(x_n) + \varphi(x_{n+1})$  and  $s_n = d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n)$ , that

$$0 \leq \limsup_{n \rightarrow +\infty} \zeta(d(x_n, x_{n+1}) + \varphi(x_n) + \varphi(x_{n+1}), d(x_{n-1}, x_n) + \varphi(x_{n-1}) + \varphi(x_n)) < 0,$$

which is a contradiction. Then we conclude that  $r = 0$  and from (3.3), since  $\varphi \geq 0$ , we get

$$\lim_{n \rightarrow +\infty} d(x_{n-1}, x_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} \varphi(x_n) = 0.$$

The second step is to prove that the sequence  $\{x_n\}$  is bounded. Let us assume that  $\{x_n\}$  is not a bounded sequence. Then, there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $n_1 = 1$  and for each  $k \in \mathbb{N}$ ,  $n_{k+1}$  is the minimum integer greater than  $n_k$  such that

$$d(x_{n_{k+1}}, x_{n_k}) > 1$$

and

$$d(x_m, x_{n_k}) \leq 1, \text{ for } n_k \leq m \leq n_{k+1} - 1.$$

By the triangle inequality, we obtain

$$\begin{aligned} 1 < d(x_{n_{k+1}}, x_{n_k}) &\leq d(x_{n_{k+1}}, x_{n_{k+1}-1}) + d(x_{n_{k+1}-1}, x_{n_k}) \\ &\leq d(x_{n_{k+1}}, x_{n_{k+1}-1}) + 1. \end{aligned}$$

Letting  $k \rightarrow +\infty$  in the above inequality and using (3.2), we get

$$\lim_{k \rightarrow +\infty} d(x_{n_{k+1}}, x_{n_k}) = 1. \tag{3.4}$$

From

$$1 < d(x_{n_{k+1}}, x_{n_k}) \leq d(x_{n_{k+1}}, x_{n_{k+1}-1}) + d(x_{n_{k+1}-1}, x_{n_k-1}) + d(x_{n_k-1}, x_{n_k})$$

and (3.2), we deduce that we can assume  $d(x_{n_{k+1}-1}, x_{n_k-1}) > 0$  for all  $k \in \mathbb{N}$ .

Again, from (3.1) and  $(\zeta_2)$ , we deduce

$$\begin{aligned} 1 < d(x_{n_{k+1}}, x_{n_k}) &\leq d(x_{n_{k+1}}, x_{n_k}) + \varphi(x_{n_{k+1}}) + \varphi(x_{n_k}) \\ &< d(x_{n_{k+1}-1}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1}) \\ &\leq d(x_{n_{k+1}-1}, x_{n_k}) + d(x_{n_k}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1}) \\ &\leq 1 + d(x_{n_k}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1}). \end{aligned}$$

Letting  $k \rightarrow +\infty$  in the above inequality, using (3.2), we get

$$\begin{aligned} \lim_{k \rightarrow +\infty} d(x_{n_{k+1}}, x_{n_k}) + \varphi(x_{n_{k+1}}) + \varphi(x_{n_k}) &= 1, \\ \lim_{k \rightarrow +\infty} d(x_{n_{k+1}-1}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1}) &= 1. \end{aligned}$$

Then by condition  $(\zeta_3)$ , with  $t_k = d(x_{n_{k+1}}, x_{n_k}) + \varphi(x_{n_{k+1}}) + \varphi(x_{n_k})$  and  $s_k = d(x_{n_{k+1}-1}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1})$ , we obtain

$$0 \leq \limsup_{k \rightarrow +\infty} \zeta \left( d(x_{n_{k+1}}, x_{n_k}) + \varphi(x_{n_{k+1}}) + \varphi(x_{n_k}), d(x_{n_{k+1}-1}, x_{n_k-1}) + \varphi(x_{n_{k+1}-1}) + \varphi(x_{n_k-1}) \right) < 0,$$

which is a contradiction. Thus the sequence  $\{x_n\}$  is bounded.

The third step is to prove that the sequence  $\{x_n\}$  is Cauchy. To this end, let

$$C_n = \sup\{d(x_i, x_j) : i, j \geq n\}, \quad n \in \mathbb{N}.$$

Since  $\{x_n\}$  is bounded, we know that  $C_n < +\infty$  for every  $n \in \mathbb{N}$ . Note that  $\{C_n\}$  is a positive non-increasing sequence and hence there exists some  $C \geq 0$  such that

$$\lim_{n \rightarrow +\infty} C_n = C. \tag{3.5}$$

Let us suppose that  $C > 0$ . By the definition of  $C_n$ , for every  $k \in \mathbb{N}$ , there exist  $n_k, m_k \in \mathbb{N}$  such that  $m_k > n_k \geq k$  and

$$C_k - \frac{1}{k} < d(x_{m_k}, x_{n_k}) \leq C_k.$$

Letting  $k \rightarrow +\infty$  in the above inequality, we get

$$\lim_{k \rightarrow +\infty} d(x_{m_k}, x_{n_k}) = C. \tag{3.6}$$

By (3.6), we can assume that  $d(x_{m_k}, x_{n_k}) > 0$  for all  $k \in \mathbb{N}$ . From

$$d(x_{m_k}, x_{n_k}) \leq d(x_{m_k}, x_{m_k-1}) + d(x_{m_k-1}, x_{n_k-1}) + d(x_{n_k-1}, x_{n_k}),$$

(3.2) and (3.6), we deduce that we can assume  $d(x_{m_k-1}, x_{n_k-1}) > 0$  for all  $k \in \mathbb{N}$ .

Again, from (3.1), (3.2), ( $\zeta_2$ ) and the definition of  $C_n$ , we deduce

$$\begin{aligned} d(x_{m_k}, x_{n_k}) &\leq d(x_{m_k}, x_{n_k}) + \varphi(x_{m_k}) + \varphi(x_{n_k}) \\ &\leq d(x_{m_k-1}, x_{n_k-1}) + \varphi(x_{m_k-1}) + \varphi(x_{n_k-1}) \\ &\leq C_{k-1} + \varphi(x_{m_k-1}) + \varphi(x_{n_k-1}). \end{aligned}$$

Letting  $k \rightarrow +\infty$  in the above inequality, using (3.2), (3.5) and (3.6), we obtain

$$\begin{aligned} \lim_{k \rightarrow +\infty} d(x_{m_k}, x_{n_k}) + \varphi(x_{m_k}) + \varphi(x_{n_k}) &= C, \\ \lim_{k \rightarrow +\infty} d(x_{m_k-1}, x_{n_k-1}) + \varphi(x_{m_k-1}) + \varphi(x_{n_k-1}) &= C. \end{aligned}$$

By condition ( $\zeta_3$ ), with  $t_k = d(x_{m_k}, x_{n_k}) + \varphi(x_{m_k}) + \varphi(x_{n_k})$  and  $s_k = d(x_{m_k-1}, x_{n_k-1}) + \varphi(x_{m_k-1}) + \varphi(x_{n_k-1})$ , we get

$$0 \leq \limsup_{k \rightarrow +\infty} \zeta(d(x_{m_k}, x_{n_k}) + \varphi(x_{m_k}) + \varphi(x_{n_k}), d(x_{m_k-1}, x_{n_k-1}) + \varphi(x_{m_k-1}) + \varphi(x_{n_k-1})) < 0,$$

which is a contradiction. Thus we have  $C = 0$  and this ensures that  $\{x_n\}$  is a Cauchy sequence. □

Now, we present our first main result inspired to the papers [10, 26, 28].

**Theorem 3.2.** *Let  $(X, d)$  be a complete metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist a simulation function  $\zeta$  and a lower semi-continuous function  $\varphi : X \rightarrow [0, +\infty[$  such that (3.1) holds, that is,*

$$\zeta(d(Tx, Ty) + \varphi(Tx) + \varphi(Ty), d(x, y) + \varphi(x) + \varphi(y)) \geq 0 \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point  $z$  such that  $\varphi(z) = 0$ .*

*Proof.* Let  $x_0 \in X$  and  $\{x_n\}$  be a sequence of Picard with initial point at  $x_0$ . At first, observe that if  $x_m = x_{m+1}$  for some  $m \in \mathbb{N}$ , then  $x_m = x_{m+1} = Tx_m$ , that is,  $x_m$  is a fixed point of  $T$ . In this case, the existence of a fixed point is proved. Therefore, we can suppose that  $x_n \neq x_{n+1}$  for every  $n \in \mathbb{N}$ .

Now, by Lemma 3.1, the sequence  $\{x_n\}$  is Cauchy and since  $(X, d)$  is complete, then there exists some  $z \in X$  such that

$$\lim_{n \rightarrow +\infty} x_n = z. \tag{3.7}$$

Note that first step of Lemma 3.1 and lower semi-continuity of the function  $\varphi$  give

$$0 \leq \varphi(z) \leq \liminf_{n \rightarrow +\infty} \varphi(x_n) = 0,$$

that is,  $\varphi(z) = 0$ .

We claim that  $z$  is a fixed point of  $T$ . If there exists a subsequence  $x_{n_k}$  of  $x_n$  such that  $x_{n_k} = z$  or  $Tx_{n_k} = Tz$  for all  $k \in \mathbb{N}$ , then  $z$  is a fixed point for  $T$ . If this does not occur, then we can assume that  $x_n \neq z$  and  $Tx_n \neq Tz$  for all  $n \in \mathbb{N}$ . Using (3.1) and  $(\zeta_2)$  with  $x = x_n$  and  $y = z$ , we deduce that

$$\begin{aligned} 0 &\leq \zeta(d(Tx_n, Tz) + \varphi(Tx_n) + \varphi(Tz), d(x_n, z) + \varphi(x_n) + \varphi(z)) \\ &< d(x_n, z) + \varphi(x_n) + \varphi(z) - [d(Tx_n, Tz) + \varphi(Tx_n) + \varphi(Tz)]. \end{aligned}$$

This implies

$$d(Tx_n, Tz) + \varphi(Tx_n) + \varphi(Tz) < d(x_n, z) + \varphi(x_n) + \varphi(z) \quad \text{for all } n \in \mathbb{N}$$

and consequently

$$\begin{aligned} d(z, Tz) &\leq d(z, x_{n+1}) + d(Tx_n, Tz) \\ &\leq d(z, x_{n+1}) + d(Tx_n, Tz) + \varphi(Tx_n) + \varphi(Tz) \\ &< d(z, x_{n+1}) + d(x_n, z) + \varphi(x_n) + \varphi(z) \end{aligned}$$

for all  $n \in \mathbb{N}$ . Letting  $n \rightarrow +\infty$  in the above inequality, we obtain that  $d(z, Tz) = 0$ , that is,  $z = Tz$ .

Now, we establish uniqueness of the fixed point. Suppose that there exists  $w \in X$  such that  $w = Tw$  and  $z \neq w$ . Using (3.1) and  $(\zeta_2)$  with  $x = w$  and  $y = z$ , we get that

$$0 \leq \zeta(d(fw, fz) + \varphi(w) + \varphi(z), d(w, z) + \varphi(w) + \varphi(z)) < d(w, z) - d(w, z) = 0,$$

which is a contradiction and hence  $w = z$ . This ends the proof of Theorem 3.2. □

#### 4. Consequences

We show the unifying power of simulation functions by applying Theorem 3.2 to deduce different kinds of contractive conditions in the existing literature.

The following corollary is a result of Banach type [2].

**Corollary 4.1.** *Let  $(X, d)$  be a complete metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist  $\lambda \in ]0, 1[$  and a lower semi-continuous function  $\varphi : X \rightarrow [0, +\infty[$  such that*

$$d(Tx, Ty) + \varphi(Tx) + \varphi(Ty) \leq \lambda d(x, y) + \varphi(x) + \varphi(y) \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point.*

*Proof.* The result follows from Theorem 3.2, by taking as simulation function

$$\zeta(t, s) = \lambda s - t,$$

for all  $t, s \geq 0$ . □

Note that we obtain Banach contraction principle if we assume  $\varphi(x) = 0$  for all  $x \in X$ .

The following corollary is a result of Rhoades type [22].

**Corollary 4.2.** *Let  $(X, d)$  be a complete metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist two lower semi-continuous functions  $\psi : [0, +\infty[ \rightarrow [0, +\infty[$  with  $\psi^{-1}(0) = \{0\}$  and  $\varphi : X \rightarrow [0, +\infty[$  such that*

$$d(Tx, Ty) + \varphi(Tx) + \varphi(Ty) \leq d(x, y) + \varphi(x) + \varphi(y) - \psi(d(x, y) + \varphi(x) + \varphi(y)) \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point.*

*Proof.* The result follows from Theorem 3.2, by taking as simulation function

$$\zeta(t, s) = s - \psi(s) - t,$$

for all  $t, s \geq 0$ . □

We have also the following corollary, see [19].

**Corollary 4.3.** *Let  $(X, d)$  be a complete metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist a function  $\psi : [0, +\infty[ \rightarrow [0, 1[$  with  $\limsup_{t \rightarrow r^+} \psi(t) < 1$  for all  $r > 0$  and a lower semi-continuous function  $\varphi : X \rightarrow [0, +\infty[$  such that*

$$d(Tx, Ty) + \varphi(Tx) + \varphi(Ty) \leq \psi(d(x, y) + \varphi(x) + \varphi(y)) [d(x, y) + \varphi(x) + \varphi(y)] \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point.*

*Proof.* The result follows from Theorem 3.2, by taking as simulation function

$$\zeta(t, s) = s\psi(s) - t,$$

for all  $t, s \geq 0$ . □

The following corollary is a result of Boyd-Wong type [4].

**Corollary 4.4.** *Let  $(X, d)$  be a complete metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exist an upper semi-continuous function  $\eta : [0, +\infty[ \rightarrow [0, +\infty[$  with  $\eta(t) < t$  for all  $t > 0$  and  $\eta(0) = 0$  and a lower semi-continuous function  $\varphi : X \rightarrow [0, +\infty[$  such that*

$$d(Tx, Ty) + \varphi(Tx) + \varphi(Ty) \leq \eta(d(x, y) + \varphi(x) + \varphi(y)) \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point.*

*Proof.* The result follows from Theorem 3.2, by taking as simulation function

$$\zeta(t, s) = \eta(s) - t,$$

for all  $t, s \geq 0$ . □

Note that we obtain Boyd-Wong result if we assume  $\varphi(x) = 0$  for all  $x \in X$ .

Next example shows that Theorem 3.2 is a proper generalization of both Banach contraction principle and Boyd-Wong result in the setting of metric spaces.

**Example 4.5** ([28], Example 4). Let  $X = [0, 1]$  endowed with the usual metric  $d(x, y) = |x - y|$  for all  $x, y \in X$ . Obviously  $(X, d)$  is a complete metric space. Also, fix  $r \in [0, 1[$ , and define  $T : X \rightarrow X$  by

$$Tx = \begin{cases} 0 & \text{if } x = 0, \\ \frac{r}{2n} - r \frac{2n-1}{2n} (2nx - 1) & \text{if } \frac{1}{2n} \leq x \leq \frac{1}{2n-1}, \\ \frac{r}{2n} + r \frac{2n+1}{2n} (2nx - 1) & \text{if } \frac{1}{2n+1} \leq x \leq \frac{1}{2n}. \end{cases}$$

Firstly, we prove that  $T$  is not a contraction. In fact, if for odd  $n > 1$  we choose  $x = \frac{1}{2n-1}$  and  $y = \frac{1}{n-1}$ , we have

$$d(Tx, Ty) = \frac{r}{n-1} \leq d(x, y) = \frac{n}{(n-1)(2n-1)} \leq \frac{3}{5(n-1)},$$



which is not satisfied for  $r > 3/5$ . Therefore, both the Banach contraction principle and Boyd-Wong result cannot be applied.

On the other hand, if we consider the function  $\phi : X \rightarrow [0, +\infty[$  defined by  $\phi(x) = x$ , then we obtain

$$\begin{aligned} d(Tx, Ty) + \phi(Tx) + \phi(Ty) &= 2 \max\{Tx, Ty\} \\ &\leq 2 \max\{rx, ry\} \\ &= r2 \max\{x, y\} \\ &= r[d(x, y) + \phi(x) + \phi(y)], \end{aligned}$$

for all  $x, y \in X$ . Thus, all the conditions of Corollary 4.1 are satisfied. Therefore,  $T$  has a unique fixed point in  $X$ .

### 5. Fixed points in partial metric spaces

In this section, from our Theorem 3.2, we deduce easily various fixed point theorems on partial metric spaces including the Matthews fixed point theorem.

**Theorem 5.1.** *Let  $(X, p)$  be a complete partial metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exists a simulation function  $\zeta$  such that*

$$\zeta(p(Tx, Ty), p(x, y)) \geq 0 \quad \text{for all } x, y \in X. \tag{5.1}$$

Then  $T$  has a unique fixed point  $z \in X$  such that  $p(z, z) = 0$ .

*Proof.* From (2.1), we deduce that

$$p(x, y) = \frac{p^s(x, y) + p(x, x) + p(y, y)}{2} \quad \text{for all } x, y \in X. \tag{5.2}$$

By Lemma 2.2, the metric space  $(X, 2^{-1}p^s)$  is complete since  $(X, p)$  is complete. Also, by Lemma 2.1, the function  $\varphi : X \rightarrow [0, +\infty[$  defined by  $\varphi(x) = 2^{-1}p(x, x)$  is lower semi-continuous in  $(X, 2^{-1}p^s)$ . Now, from (5.1) and (5.2), we obtain that the mapping  $T$  satisfies the following contractive condition

$$\zeta(2^{-1}p^s(Tx, Ty) + \varphi(Tx) + \varphi(Ty), 2^{-1}p^s(x, y) + \varphi(x) + \varphi(y)) \geq 0 \quad \text{for all } x, y \in X.$$

Thus the mapping  $T$  satisfies all the condition of Theorem 3.2 with respect to the metric space  $(X, 2^{-1}p^s)$ , and hence has a unique fixed point  $z \in X$  such that  $p(z, z) = 2\varphi(z) = 0$ . □

Now, the Matthews fixed point theorem follows immediately from Theorem 5.1 by choosing as simulation function

$$\zeta(t, s) = \lambda s - t \quad \text{for all } t, s \in [0, +\infty[$$

with  $\lambda \in [0, 1[$ .

**Corollary 5.2.** *Let  $(X, p)$  be a complete partial metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exists a  $\lambda \in [0, 1[$  such that*

$$p(Tx, Ty) \leq \lambda p(x, y) \quad \text{for all } x, y \in X. \tag{5.3}$$

Then  $T$  has a unique fixed point  $z \in X$  such that  $p(z, z) = 0$ .

If we choose as simulation function

$$\zeta(t, s) = \eta(s) - t \quad \text{for all } t, s \in [0, +\infty[$$

where  $\eta : [0, +\infty[ \rightarrow [0, +\infty[$  is an upper semi-continuous function with  $\eta(t) < t$  for all  $t > 0$  and  $\eta(0) = 0$ , then we obtain the following result of Boyd-Wong type in the setting of partial metric space.

**Corollary 5.3.** *Let  $(X, p)$  be a complete partial metric space and let  $T : X \rightarrow X$  be a mapping. Suppose that there exists an upper semi-continuous function  $\eta : [0, +\infty[ \rightarrow [0, +\infty[$  with  $\eta(t) < t$  for all  $t > 0$  and  $\eta(0) = 0$  such that*

$$d(Tx, Ty) \leq \eta(d(x, y)) \quad \text{for all } x, y \in X.$$

*Then  $T$  has a unique fixed point.*

Appropriately choosing the simulation function we can obtain other known results of fixed point in partial metric spaces.

## References

- [1] H. Argoubi, B. Samet, C. Vetro, *Nonlinear contractions involving simulation functions in a metric space with a partial order*, (submitted). 1, 2, 2.6
- [2] S. Banach, *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fund. Math., **3** (1922), 133–181.1, 4
- [3] V. Berinde, *Generalized contractions in quasimetric spaces*, Seminar on Fixed Point Theory, Babe-Bolyai Univ. Cluj-Napoca, **3** (1993), 3–9.1
- [4] D. W. Boyd, J. S. W. Wong, *On nonlinear contractions*, Proc. Amer. Math. Soc., **20** (1969), 458–464.4
- [5] M. A. Bukatin, J. S. Scott, *Towards computing distances between programs via Scott domains*, Logical Foundations of Computer Science, Lecture Notes in Comput. Sci. Springer Berlin, **1234** (1997), 33–43.1
- [6] M. A. Bukatin, S. Y. Shorina, *Partial metrics and co-continuous valuations*, Foundations of Software Science and Computation Structures, Lecture Notes in Computer Science (ed. M. Nivat), Springer Berlin, **1378** (1998), 33–43.1
- [7] L. Ćirić, B. Samet, C. Vetro, *Common fixed point theorems for families of occasionally weakly compatible mappings*, Math. Comput. Modelling, **53** (2011), 631–636.1
- [8] L. Ćirić, B. Samet, C. Vetro, M. Abbas, *Fixed point results for weak contractive mappings in ordered  $K$ -metric spaces*, Fixed Point Theory, **13** (2012), 59–72.1
- [9] M. Cosentino, P. Salimi, P. Vetro, *Fixed point results on metric-type spaces*, Acta Math. Sci. Ser. B Engl. Ed., **34** (2014), 1237–1253.1
- [10] F. Khojasteh, S. Shukla, S. Radenović, *A new approach to the study of fixed point theorems via simulation functions*, Filomat, (2014), in press. 1, 2, 2.4, 3
- [11] V. Lakshmikantham, L. Ćirić, *Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces*, Nonlinear Anal., **70** (2009), 4341–4349.1
- [12] S. G. Matthews, *Partial metric topology*, Proc. 8th Summer Conference on General Topology and Applications, Ann. New York Acad. Sci., **728** (1994), 183–197.1, 2, 2.2
- [13] S. J. O’Neill, *Partial metrics, valuations and domain theory*, Conference on General Topology and Applications, Ann. New York Acad. Sci., **806** (1996), 304–315.1, 2
- [14] S. Oltra, O. Valero, *Banach’s fixed point theorem for partial metric spaces*, Rend. Istit. Mat. Univ. Trieste, **36** (2004), 17–26.2, 2.2
- [15] D. Paesano, P. Vetro, *Suzuki’s type characterizations of completeness for partial metric spaces and fixed points for partially ordered metric spaces*, Topology Appl., **159** (2012), 911–920.2
- [16] S. Romaguera, *A Kirk type characterization of completeness for partial metric spaces*, Fixed Point Theory Appl., **2010** (2010), 6 pages.2
- [17] S. Romaguera, M. Schellekens, *Partial metric monoids and semivaluation spaces*, Topology Appl., **153** (2005), 948–962.1
- [18] D. Reem, S. Reich, A. J. Zaslavski, *Two Results in Metric Fixed Point Theory*, J. Fixed Point Theory Appl., **1** (2007), 149–157.1
- [19] S. Reich, *Fixed points of contractive functions*, Boll. Un. Mat. Ital., **5** (1972), 26–42.4
- [20] S. Reich, A. J. Zaslavski, *A Fixed Point Theorem for Matkowski Contractions*, Fixed Point Theory, **8** (2007), 303–307.1
- [21] S. Reich, A. J. Zaslavski, *A Note on Rakotch contraction*, Fixed Point Theory, **9** (2008), 267–273.1
- [22] B. E. Rhoades, *Some theorems on weakly contractive maps*, Nonlinear Anal., **47** (2001), 2683–2693.4
- [23] A. Roldán, E. Karapinar, C. Roldán, J. Martínez-Moreno, *Coincidence point theorems on metric spaces via simulation functions*, J. Comput. Appl. Math., **275** (2015), 345–355.1
- [24] I. A. Rus, *Generalized Contractions and Applications*, Cluj University Press, Cluj-Napoca, (2001).1
- [25] I. A. Rus, A. Petruşel, G. Petruşel, *Fixed Point Theory*, Cluj University Press, Cluj-Napoca, (2008).1
- [26] B. Samet, C. Vetro, F. Vetro, *From metric spaces to partial metric spaces*, Fixed Point Theory Appl., **2013** (2013), 11 pp.1, 3
- [27] C. Vetro, F. Vetro, *Common fixed points of mappings satisfying implicit relations in partial metric spaces*, J. Nonlinear Sci. Appl., **6** (2013), 152–161.1

- [28] C. Vetro, F. Vetro, *Metric or partial metric spaces endowed with a finite number of graphs: a tool to obtain fixed point results*, Topology Appl., **164** (2014), 125–137.1, 3, 4.5